

제 81회

ORGAN ON A CHIP

기술교류회

2022.04.14 목 오후 4시 30분

한림대학교 자연과학관 7103호



황재성 교수

경희대학교

1. Education

박사: 아주대학교 의과대학 (2003)

석사: 서울대학교 (1994)

학사: 서울대학교 (1992)

2. Experience

2020 ~ 현재

2009 ~ 현재

1994 ~ 2009

피부과학 응용소재 선도기술개발 사업단, 단장
경희대학교 생명과학대학 유전생명공학과, 교수
(주)아모레퍼시픽 피부연구팀, 피부의약연구팀 팀장,
수석연구원

제목

멜라닌 형성세포에서 멜라닌 이동 조절 기전

Regulation of skin pigmentation by interrupting melanosome transport and finding a novel function of Prohibitin on it

초록

Melanocytes are unique cells that produce specific melanin-containing intracellular organelles called melanosomes. Melanosomes are lysosome-related organelles specialized in melanin synthesis. The movement of melanosome was mediated by the formation of a Rab27a-Melanophilin-MyosinVa protein complex. Melanosomes are transported from the perinuclear area of melanocytes toward the plasma membrane as they become more melanized and finally transferred to keratinocyte. We screened several compounds which can affect melanosome transport in melanocytes. Using a pharmacophore modeling, we found a compound which inhibits the binding of Rab27a and Melanophilin. We also found some compounds which affect the expression of Rab27a, Melanophilin and MyosinVa from a cell-based screening. We showed that the regulation of these proteins can affect skin pigmentation from reconstituted skin or animal study. These results showed that regulation of melanosome transport can be a good target for searching depigmenting agent.

Prohibitin (PHB, also known as PHB1 or BAP32), is a highly conserved 31kDa protein that expressed in many cellular compartments, such as mitochondria, nucleus, cytosol, and plasma membrane, and plays roles in regulating the transcription of genes, apoptosis, and mitochondrial biogenesis. There is a report that Prohibitin expression is required for the stimulation of pigmentation by melanogenin. However, no studies have been published on the function of PHB in melanocytes, especially in melanosome transport. We found that PHB is located in the melanosome and perinuclear aggregation of melanosome is induced when expression of PHB is reduced with no influence on melanin contents. PHB binds directly to Rab27a and Mlph but not Myosin-Va. Rab27a and Mlph bind to specific domains of PHB. Reduced expression of PHB led to the impaired binding affinity between Rab27a and Mlph. From these results, we found that PHB regulates melanosome transport by linking to Rab27a and Mlph in melanocytes. Targeting and regulating PHB not only manages pigmentation in melanocytes, but also controls hyperpigmentation in melanoma.

주 관 한림대학교 미래융합스쿨 융합신소재공학전공, 융합신소재공학연구소

후 원 한국연구재단 중견연구사업, 산업통상자원부 3D 생체조직칩 제품화사업

지 원 한림대학교 대학원 나노-메디컬 디바이스 공학 협동과정, 춘천바이오산업진흥원

문의처: de3553@hallym.ac.kr / Tel: 033-248-3553